



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

Improved functional and histochemical outcomes in l-DOPA plus tolcapone treated VMAT2-deficient mice

Moreira, Carlos G ; Morawska, Marta M ; Baumann, Aron ; Masneuf, Sophie ; Linnebank, Michael ; Sommerauer, Michael ; Landolt, Hans-Peter ; Noain, Daniela ; Baumann, Christian R

Abstract: Parkinson disease is typically treated with L-3,4-dihydroxyphenylalanine (or levodopa) co-prescribed with concentration stabilizers to prevent undesired motor fluctuations. However, the beneficial role of the chronic combined therapy on disease progression has not been thoroughly explored. We hypothesized that tolcapone, a catechol-O-methyl-transferase inhibitor, co-administered with levodopa may offer beneficial long-term disease-modifying effects through its dopamine stabilization actions. Here, we followed vesicular monoamine transporter 2-deficient and wild-type mice treated twice daily per os with vehicle, levodopa (20 mg/kg), tolcapone (15 mg/kg) or levodopa (12.5 mg/kg) + tolcapone (15 mg/kg) for 17 weeks. We assessed open field, bar test and rotarod performances at baseline and every 4th week thereafter, corresponding to OFF-medication weeks. Finally, we collected coronal sections from the frontal caudate-putamen and determined the reactivity level of dopamine transporter. Vesicular monoamine transporter 2-deficient mice responded positively to chronic levodopa + tolcapone intervention in the bar test during OFF-periods. Neither levodopa nor tolcapone interventions offered significant improvements on their own. Similarly, chronic levodopa + tolcapone intervention was associated with partially rescued dopamine transporter levels, whereas animals treated solely with levodopa or tolcapone did not present this effect. Interestingly, 4-month progression of bar test scores correlated significantly with dopamine-transporter-label density. Overall, we observed a moderate functional and histopathological improvement effect by chronic dopamine replacement when combined with tolcapone in vesicular monoamine transporter 2-deficient mice. Altogether, chronic stabilization of dopamine levels by catechol-O-methyl-transferase inhibition, besides its intended immediate actions, arises as a potential long-term beneficial approach during the progression of Parkinson disease.

DOI: <https://doi.org/10.1016/j.neuropharm.2020.108353>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-191449>

Journal Article

Accepted Version

Originally published at:

Moreira, Carlos G; Morawska, Marta M; Baumann, Aron; Masneuf, Sophie; Linnebank, Michael; Sommerauer, Michael; Landolt, Hans-Peter; Noain, Daniela; Baumann, Christian R (2020). Improved functional and histochemical outcomes in l-DOPA plus tolcapone treated VMAT2-deficient mice. *Neuropharmacology*, 181:108353.

DOI: <https://doi.org/10.1016/j.neuropharm.2020.108353>

1. Introduction (we don't have a title and an abstract page?)

Oral administration of L-3,4-dihydroxyphenylalanine (levodopa or L-DOPA), the endogenous precursor of dopamine (DA), combined with a DA decarboxylase inhibitor, such as benserazide hydrochloride or carbidopa, is currently the most effective, tolerable, and therefore broadly used treatment for patients with Parkinson disease (PD). However, the use of L-DOPA is burdened with several limitations, including short-time bioavailability of each dose due to extensive peripheral metabolism. In addition, there is the assumption that long-term dopaminergic stimulation with L-DOPA facilitates the occurrence of motor fluctuations with dyskinesia (Connolly and Lang, 2014; Jenner et al., 2011). The latter have been closely related to disease duration, and treatment duration and dosage (Schrag and Quinn, 2000).

Recent recommendations, therefore, suggested to evaluate alternative pharmacotherapies than L-DOPA for treatment initiation in certain patients, to avoid long-term levodopa-related motor complications (Connolly and Lang, 2014). On the other hand, an elegant delayed-start trial in early PD patients recently found similar rates of dyskinesia and L-DOPA-related motor fluctuations at 80 weeks when comparing a group treated with L-DOPA for 80 weeks and another group placebo-treated for 40 weeks followed by 40 weeks of L-DOPA treatments (Verschuur et al., 2019). The authors concluded that L-DOPA treatment duration may not havebear a disease-modifying effect, at least in relation to motor fluctuations.

In addition to treatment duration and dosing, the pattern of DA receptor stimulation has been associated with the emergence of dyskinesias: continuous dopaminergic treatment - either with long-acting DA-receptor agonists or continuous L-DOPA

infusions - is related to fewer ~~motor fluctuations during dyskinesias and motor~~
~~fluctuations, as for instance evidenced in a primate PD model on medication~~ (Bibbiani
et al., 2005). This finding has driven the search for strategies to reduce pulsatile DA
receptor stimulation (Nyholm, 2006; Olanow and Schapira, 2013; Chondrogiorgi et al.,
2014). The goal of the STRIDE-PD study was to examine whether adding entacapone,
an inhibitor of catechol-O-methyltransferase (COMT), to L-DOPA reduces the
development of dyskinesia, as this approach increases L-DOPA's half-life and
smoothen its serum concentrations, i.e. leads to a mildly enhanced continuous
stimulation of postsynaptic DA receptors (Stocchi and Olanow, 2004; Stocchi et al.,
2010). The study was negative, but the authors rightfully discussed the outcome's
relation with the study design, as it provided higher L-DOPA dose equivalents in the
entacapone group. ~~In fact~~ On the other hand, experimental animal models of LIDs (L-
DOPA induced dyskinesias) suggest that the loss of nigrostriatal DA neurons may be
the cause for or at least contribute to the development of such abnormal involuntary
movements (Cenci, 2014). ~~All in all~~ Altogether, despite the extensive and negative
STRIDE study, it is still unclear whether COMT inhibition during prolonged long-term
dopaminergic stimulation at bioavailable L-DOPA equivalent doses offers any long-
lasting and/or disease-modifying effect in PD, such as which would be best reflected
by deceleration of nigrostriatal DA denervation, and which in turn, might have a
beneficial impact on LIDs onset and severity.

Tolcapone, a selective and reversible COMT inhibitor penetrating the blood-brain
barrier (Gulberg and Marsden, 1975; Backstrom et al., 1989; Kopin, 1985; Nagatsu
and Sawada, 2009a; Kiss et al., 2010), offers a potent and sustained inhibition *in vivo*
(Zürcher et al., 1990; Bonifácio et al., 2007). ~~It is widely accepted that central COMT~~
~~inhibition by tolcapone reduces degradation of L-DOPA to 3-O-methyldopa, thus~~

widening its therapeutic window, while further prolonging L-DOPA effectiveness in the brain by reducing DA metabolism to homovanillic acid (Kurth and Adler, 1998; Kaakkola, 2000; Zeuner et al., 2019). Besides peripherally inhibiting L-DOPA metabolism, tolcapone further inactivates central COMT and reduces degradation of extrasynaptic or diffused DA to 3-methoxytyramine (Kurth and Adler, 1998; Kaakkola, 2000; Zeuner et al., 2019), thus, stabilizing its bioavailable synaptic concentration and modulating dopaminergic transmission of presynaptic and postsynaptic neurons (Chen et al., 2011). Ultimately, continuous postsynaptic dopaminergic signaling by tempered and stable DA levels will possibly lead to long-term clinical and pathological amelioration. Notwithstanding, tolcapone was associated with increases in liver enzymes in a dose-dependent manner (Longo et al., 2016). Peripheral inhibition of soluble-COMT may be at least in part, the main underlying cause of tolcapone-induced hepatotoxicity: uncoupling of oxidative phosphorylation and the subsequent reduction in mitochondrial energy (Haasio et al., 2002).

The cause of L-DOPA-induced motor fluctuations is not entirely understood. Apart from contributions from disturbed neurotransmission, the best available evidence suggests that progressive degeneration of DA-producing cells in the substantia nigra is mandatory (Leta et al., 2019). In parallel, degeneration of dopaminergic terminals along with aberrant processing of exogenously administered L-DOPA lead to imbalanced striatal DA levels, causing motor fluctuations (Niccolini et al., 2015). For instance, the link of dyskinesia with dopaminergic denervation is supported by findings in 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) addicts and in rodents with complete nigral lesions (Winkler et al., 2002; Langston, 2017), although these studies lacked a necessary longitudinal perspective. Altogether, there appears to be a strong

~~association between the degree of nigrostriatal denervation/neurodegeneration and the expression of dyskinesia (Di Monte et al., 2000).~~

~~Ultimately contributing to neurodegeneration as well, chronic~~Chronic use of L-DOPA generates several toxic molecules ~~(Paul and Borah, 2016),~~ responsible for increased predisposition to ~~vascular disease and~~ accelerated disease progression (Müller, 2011). One such metabolite is homocysteine (Hcy), the level of which increases in the plasma of PD patients ~~ultimately generating~~composes a high risk for vascular diseases and dementia (Kocer et al., 2016). Furthermore, Hcy exerts a neurotoxic action and may participate in the mechanisms of neurodegeneration, such as DNA damage, excitotoxicity, oxidative stress ~~and neuronal degeneration (Bhattacharjee and Borah,~~ calcium accumulation, and apoptosis (Zoccolella et al., 2006a). Several studies have shown that higher concentration of Hcy in PD is related to long-term administration of L-DOPA (Lamberti et al., 2005; Ibrahimagic et al., 2016). Therapeutic approaches against ~~hyperhomocysteinemia~~hyperhomocysteinaemia include COMT inhibition, reducing S-adenosylhomocysteine by blocking COMT's O-methylation of L-DOPA, and in turn lowering its subproduct, ~~Hcy~~ Hcy, and the risk of associated co-morbidities and neurodegenerative burden (Paul and Borah, 2016); ~~Zoccolella et al., 2006b).~~

To investigate COMT inhibition in DA replacement therapy in a preclinical setting, we used avesicular monoamine transporter 2-deficient (VMAT2-LO) mice. This genetic animal model of PD ~~that~~ encompasses many of the progressive motor and non-motor symptoms described in patients, particularly in aged subjects. During our window of pharmacological intervention, from 14 to 18 months of age, these animals show gradually decreased motor function, as well as ~~the hallmark neuropathology of the disease, such as the vesicular monoamine transporter 2-deficient (VMAT2-LO) mice~~ age-dependent decline of DAT expression and activity, contributing to neuronal

Formatiert: Englisch (Vereinigte Staaten), Muster: Transparent

Formatiert: Schriftart: +Textkörper (Arial)

degeneration in older animals (Taylor et al., ~~2011~~2009). We specifically aimed to examine whether adding tolcapone to L-DOPA, with consecutively mildly-enhanced prolonged dopaminergic stimulation at bioavailable L-DOPA equivalent doses, reduces the progression of motor deficits during OFF-medication and striatal neuropathology – which is reportedly ~~linked~~the cause for the development to dyskinesias ~~—and motor deficits progression.~~ In addition, as a secondary outcome, we ~~examine~~examined whether ~~adding this COMT inhibitor~~combining tolcapone with prolonged L-DOPA treatment indeed reduces plasmatic levels of total Hcy (tHcy) ~~—~~ and, possibly, contributes to an overall ameliorated condition.

Formatiert: Schriftfarbe: Schwarz, Englisch (Vereinigtes Königreich), Muster: Transparent (Weiß)

Formatiert: Schriftfarbe: Schwarz, Englisch (Vereinigtes Königreich), Muster: Transparent (Weiß)

2. Materials and Methods

2.1 Animals

In total, we included 21 VMAT2-LO and 23 VMAT2-WT fourteen-month-old male mice with C57BL/6 genetic background. The VMAT2-LO line, obtained from the group of Gary Miller at Emory University, USA (Caudle et al., 2007), expresses only ~5% of the VMAT2 protein (Mooslehner et al., 2001). We single-housed all animals with food and water provided ad libitum until completion of the studies. We kept the animals on a 12-hour light/dark cycle and constant temperature (21 - 22°C) and relative humidity (30%). We weighed all mice daily and acclimated them to the experimental room before behavioral testing. We carried our studies approved by both local and federal Swiss authorities under the license n. ZH 205/2012.

2.2 Study design

122 We aimed to determine whether combined L-DOPA + tolcapone administration, in
123 comparison to single drug administration, had an OFF-medication (i.e. after drug
124 wash-out) beneficial effect on *functional*/behavioral and histological disease hallmarks.
125 Thus, we intended to initiate interventions after the onset of the majority of motor
126 symptoms (*functional* behavioral markers) and when dopamine transporter (DAT,
127 *functional* histological marker) levels were already affected, but preceding the age
128 when neurodegeneration was first reported, indicated by tyrosine hydroxylase cell loss
129 in the substantia nigra. We based the temporal window and duration of the treatments
130 on the disease dynamics reported earlier for this model (Taylor et al., 2011).

131 At 14 months of age, we began the chronic 17-week intervention with one week of
132 baseline behavioral testing followed by four identical periods consisting of 3 weeks of
133 pharmacological treatment (ON-period) plus 1 behavioral testing week (OFF-period)
134 (**Figure 1**). We administered drugs *per os* twice daily at lights-on and 8 hours later.
135 After the last behavioral testing week (17th week, OFF-period, after 5.5 days wash-out
136 of all drugs), we [obtained plasma samples to quantify liver transaminases](#), euthanized
137 the animals and collected their brains [and livers](#) for later evaluations.

139 2.3 Pharmacological interventions

140 We grouped all mice into four treatment sets: vehicle (NaCl 0.9%, '-V'); L-
141 DOPA/benserazide (20.0 mg/kg: Madopar® 125 mg, Roche Pharma AG, Switzerland,
142 '-L'), tolcapone (15.0 mg/kg: Tasmar® 100 mg, MEDA Pharma GmHB, Switzerland, '-
143 T'); and a 1:1 solution of L-DOPA/benserazide (12.5 mg/kg) and tolcapone (15.0
144 mg/kg: '-LT'). We aimed for equalized concentrations of L-DOPA/benserazide at the
145 synaptic level in both -L and -LT groups, by multiplying the ordinary L-DOPA dosage

146 (12.5 mg/kg) by 1.6, in order to achieve the same DA bioavailability in the -L group
147 (20.0 mg/kg) (Bonifácio et al., 2015). Each mouse received a total of 120 doses
148 complemented with a daily portion of wet-food (water and crushed chow) in addition
149 to the regular chow.

150

151 *2.4 Behavioral tests*

152 We performed all tests in OFF-medication weeks (start 3.5 days after the last
153 treatment episode), during the light period. We conducted all tests in 2 consecutive
154 days, following the same order and at the same time of day.

155 To assess locomotor activity, we used the open field test (Bello et al., 2011): 4 identical
156 plexiglas boxes (46x46x40 cm) and a video camera connected to an automated
157 tracking system (EthoVision XT 9.0, Noldus, Germany). We habituated animals to the
158 setup for 15 minutes prior to the testing day. During the testing trial, we allowed each
159 animal to freely explore the arena for 30 minutes while recording spontaneous
160 horizontal distance travelled.

161 We used the bar test to assess catalepsy, a syndrome often attributed to dopaminergic
162 antagonism or failure of DA neurotransmission, and interpreted to reflect bradykinesia,
163 rigidity and abnormal posture (de Ryck et al., 1980; Alvarez-Cervera et al., 2005; Noain
164 et al., 2013). Briefly, we placed the animals sitting on the working bench over their hind
165 paws and grabbing an elevated acrylic bar with their front paws. We considered a trial
166 valid when the animal remained grabbing the bar at least 2 seconds and up to 180
167 seconds.

168 To examine motor coordination, we performed a standard mouse rotarod test
169 (UgoBasile©, Italy) (Avale et al., 2004), set at constant speed. Three hours before the

170 testing session, we subjected the animals to a 10-minute training. During the testing
171 phase, we placed the animals onto the rotating treadmill until a maximum time of 180
172 seconds.

173

174 *2.5 Food intake*

175 To examine the potential influence of feeding behavior and/or body weight on
176 pharmacodynamics, we performed a modified food-intake test during the 15th week
177 (ON-period) in all animals. Briefly, we removed all dry food-pellets from the cages and,
178 for 4 days, we only presented wet-food. We weighed the cups daily and corrected the
179 results for food evaporation.

180

181 *2.6 Total homocysteine quantification*

182 To confirm whether tolcapone is indeed helpful in reducing Hcy levels during
183 prolonged L-DOPA replacement, we assayed standard ELISA determinations of ~~the~~ Hcy
184 ~~in plasma of VMAT2-WT animals~~ Hcy using a mouse homocysteine kit (Crystal Chem,
185 Catalog #80444, USA) as per manufacturer's protocol. The absorbance readings from
186 VMAT2-LO animals samples were increased up to 20-fold across all groups compared
187 to VMAT2-WT-V animals, and therefore, considered untrustworthy given the accuracy
188 limits of the commercial kit. VMAT2-LO mice have been reported to present increased
189 levels of ~~cysteine~~ free cysteinyl-DOPA and cysteinyl-DOPAC adducts, derivatives
190 of proteins the reaction of DOPA and DOPAC quinones with free cysteine (Caudle et
191 al., 2007⁷). The specificity of the commercial kit to Hcy in plasma samples also
192 containing abnormally high levels of cysteinyl isomers was not assured by the

Formatiert: Schriftart: Arial

193 [provider.](#) for which levels of tHcy could not be specifically determined in this genotype
194 (data not shown).

195

196 *2.7 Liver enzyme quantification*

197 To test whether our heaviest drug regime (-LT), as proxy of all other treatments (Longo
198 et al., 2016), negatively affected liver function in VMAT2-LO mice, we measured
199 plasma levels of aspartate transaminase (AST) and alanine transaminase (ALT), in
200 VMAT2-LO-V and VMAT2-LO-LT, by standard blood test. Two samples from the
201 VMAT2-LO-LT group were found hemolyzed at the time of analysis, and were
202 discarded by advice of the Center for Laboratory Medicine and Pathology (USZ).

203

204 *2.8 Full-body perfusion*

205 For histological analyses, we sacrificed all animals by trans-cardiac perfusion at 18
206 months of age (exsanguination using ice-cold phosphate-buffered-saline followed by
207 perfusion of freshly prepared ice-cold 4% paraformaldehyde (Sigma Aldrich).
208 Immediately after fixation, we harvested the brains, post-fixed, dehydrated, quick-froze
209 in dry ice and stored them at -80°C. Later full-brain sectioning was done at 30 µm
210 thickness in freezing-staged microtome. Due to freezing issues with the microtome,
211 we discarded two VMAT2-LO-V brains from further analysis.

212

213 *2.9 Immunohistochemistry, microscopy and color-density quantification*

214 We included every fourth coronal section from the rostral portion of the striatum.
215 Briefly, after washing the sections in tris-buffered-saline, we blocked endogenous

216 peroxidase activity by incubating sections in 2% H₂O₂ for 30 min. We incubated free-
217 floating sections with a monoclonal rat anti-DAT antibody (#MAB369; Merck; 1:500)
218 overnight at 4°C. Next, we incubated the sections in a biotinylated goat anti-rat
219 secondary antibody (#AB97057, Abcam, 1:1000) for 2 hours at RT followed by avidin-
220 biotin complex for 1.5 hours at RT. We developed sections with 0.025% 3,3'-
221 diaminobenzidine (DAB) and 0.05% H₂O₂ in tris-buffered-saline.

222 Using an Axio Imager M2 microscope (Karl Zeiss, Jena, Germany), we acquired
223 images from the coronal section with the highest signal intensity for quantification of
224 DAT stained terminals (Ruifrok and Johnston, 2001). We obtained full-section images
225 with a 2.5x objective, and further analyzed them with ImageJ (version 1.47j, National
226 Institutes of Health, USA). Briefly, we processed all images using the hematoxylin-
227 3,3'-diaminobenzidine vector, in order to obtain the DAB exclusive quantification
228 panel. From each section, the signal estimate is the average color intensity, which is
229 proportional to the concentration of the stain, of bilateral measurements in the striatum.
230 All values were corrected for area and background-noise, the last by subtracting the
231 intensity in the cortex. As per the Lambert-Beer law, we calculated DAT optical density
232 (OD, p.d.u.) according to the formula $OD = \log(255 / \text{mean_intensity_striatum}) -$
233 $\log(255 / \text{mean_intensity_cortex})$.

234

235 2.10 Statistics

236 We conducted nested 2-way ANOVA to compare the effects of all pharmacological
237 interventions on performances in open field test, bar test and rotarod, on the last OFF-
238 period of the protocol (**Figure 1**, 17th week, treatment(genotype)). Levene's test for
239 equality of variances was found to be violated for the time of immobility in the bar test

240 ($F(7,36)=3.95$, $**p=.003$), and for latency to fall in the rotarod test ($F(7,36) = 15.86$,
241 $***p<.001$), therefore data was ranked. On other instances, t-statistics not assuming
242 homogeneity of variance were corrected with Welch's test. Error bars in the figures
243 represent \pm SEM. Raw data will be made available upon reasonable request.

244

245 **3. Results**

246 *3.1 Improved progression of bradykinesia and postural stability scores OFF-* 247 *medication in VMAT2-LO mice chronically co-administered with L-DOPA and* 248 *tolcapone*

249 Determining the effect of different chronic therapeutic interventions on motor behavior
250 after washing out medication is a suitable instrument for assessing disease-modifying
251 effects. To determine the progression of motor deficits in pharmacologically intervened
252 VMAT2-LO mice and VMAT2-WT controls, we used the open field, bar and rotarod
253 tests, with a particular emphasis on so-called axial symptoms like postural
254 impairments, as such signs are considered particularly important for measuring motor
255 progression (Schreiner et al., 2019).

256 At baseline, VMAT2-LO mice performed significantly worse in all three motor tests as
257 compared to VMAT2-WT controls, that is impaired locomotor activity (unpaired
258 $t(42)=6.03$, $***p<.001$; **Figure 2a**), increased time of immobility (Welch's $t(20)=5.23$,
259 $***p<.001$; **Figure 2b**) and poorer motor coordination (unpaired $t(42)=4.17$, $***p<.001$;
260 **Figure 2c**).

261 None of the pharmacological regimes counteracted the motor deficits in VMAT2-LO
262 mice on the last assessment week, evidenced by impaired OFF-medication
263 performance compared to VMAT2-WT animals in all tests. In the open field,

collectively, VMAT2-LO showed reduced locomotor activity compared to VMAT2-WT (nested 2-way ANOVA, treatment(genotype), $F(4,36)=10.43$, $***p<.001$, **Figure 2d**) despite all treatments. For catalepsy or immobility measures, VMAT2-LO animals maintained significantly longer latency for initiation of movement than VMAT2-WT controls (nested 2-way ANOVA of ranked cases, treatment(genotype), $F(4,36)=23.06$, $***p<.001$, **Figure 2e**), however non-vehicle subjects presented slightly better marks compared to vehicle animals within each genotype. Concerning motor coordination, symptoms in VMAT2-LO remained far from VMAT2-WT in the last testing week (nested 2-way ANOVA of ranked cases, treatment(genotype), $F(4,36)=34.45$, $***p<.001$, **Figure 2f**). Adding tolcapone to L-DOPA did not significantly changed performance when compared to -L-DOPA alone or -vehicle treated VMAT2-LO animals (Tukey post hoc comparison, L*LT and V*LT, $p>.999$).

The results from the last behavioral assessment poorly reflect the longitudinal perspective of 17-weeks of progressive behavior: groups present high variability and low statistical power at the last protocol-week as well as in the other OFF-periods (5th, 9th and 13th weeks). When analyzing the global motor *progression*, from baseline to endpoint, in relation to the treatments (comparison of slopes of linear robust-fittings from each subject's scores), the progressive worsening in the bar test performance experienced by VMAT2-LO-LT mice appeared mitigated when compared to VMAT2-LO-L, (unpaired $t(8.43)=4.24$, $**p=.003$, **Figure 2k**). Alternatively, the improved performance in the bar test could be explained by prolonged efficacy of dopaminergic stimulation by the addition of tolcapone, with a clinical effect even days after the last medication intake. Apart from this isolated finding, progression of the other motor variables was similar between different treatments, again with high intra- and inter-variability (**Figure 2g-l**).

289

290 *3.2 VMAT2-LO mice co-administered with L-DOPA and tolcapone had rescued striatal*
291 *dopamine transporter levels compared to vehicle-treated transgenic mice*

292 DAT quantification in striatal tissue is a widely established marker of functional integrity
293 of the DA system and was previously used to describe disease progression in VMAT2-
294 LO mice (Caudle et al., 2007; Taylor et al., 2009). We estimated DAT reactivity in the
295 striatum (**Figure 3a**) of VMAT2-WT-V and all VMAT2-LO groups. VMAT2-LO-V mice
296 presented reduced DAT signal strength compared to VMAT2-WT-V controls (Welch's
297 $t(29.86)=2.43$, $*p=.022$, **Figure 3b**), consistent with an ongoing functional decay in
298 integrity of the striatal dopaminergic system in the mutants (Caudle et al., 2007).
299 Remarkably, VMAT2-LO-LT mice presented a partially rescued level of DAT
300 immunoreactivity ~~compared to VMAT2-LO-V (one-way(Welch's ANOVA, Brown-~~
301 ~~Forsythe $F(3,69)=4.83$, $**31.43)=9.185$, $***p=.004<.001$, **Figure 3c, d**), indicating a~~
302 ~~significantly~~ less deteriorated histopathological status ~~(Tukeycompared to VMAT2-LO-~~
303 ~~V (Games-Howell's post hoc comparison, $V*LT$, $***p=.004033$, **Figure 3d**).~~ The
304 observations between VMAT2-LO-V and VMAT2-LO-T ($p=.999=.994$), and VMAT2-
305 LO-V and VMAT2-LO-L and VMAT2-LO-LT mice ($p=.142450$) were statistically similar
306 (**Figure 3d**). VMAT2-WT are statistically equal across all groups (Kruskal-Wallis test,
307 $H(3)=.098$, $p=.992$, **Figure 3e, f**).

308

309 *3.3 Improved immobility scores are associated with rescued neuropathological status*
310 *in VMAT2-LO mice treated with L-DOPA and tolcapone*

311 There is an association between progression of motors and non-motor symptoms and
312 neuropathological status in both PD patients and animal models (Giasson et al., 2002;

313 Braak et al., 2003). To determine whether the immunoreactivity level of a functional
314 integrity striatal marker predicts well motor ability in the bar test, we calculated a
315 Pearson correlation between striatal DAT optical density and the bradykinesia
316 progression index (slope over the scores of all time points for the time of immobility in
317 the bar test) during OFF-medication in VMAT2-LO mice (**Figure 4a**). We found a
318 significant correlation (Pearson's, $r(17)=-0.50$, $*p=.015$), which could suggest an
319 association between the effect of tolcapone-mediated steadier bioavailability of L-
320 DOPA and improved motor performance progression in this model. This link was
321 visually confirmed in an exploratory group-based clustering across the correlated
322 factors, including all mutant groups and the VMAT2-WT-V group as reference (**Figure**
323 **4b**). This figure reveals that the VMAT2-LO-LT individuals (green) are closer
324 positioned to the VMAT2-WT-V (black) centroid, overall reflecting that combined
325 treatment with tolcapone provides behavioral and immunohistochemical findings that
326 are closest to those of healthy animals.

327

328 *3.4 Food intake upon different treatments in VMAT2-LO mice*

329 VMAT2-LO mice present a distinct phenotype compared to VMAT2-WT littermates,
330 thus, we characterized the feeding behavior of all animals in ON-medication. We
331 assayed a tightly-controlled protocol and found a treatment-independent food-intake
332 behavior in VMAT2-LO mice in respect to VMAT2-WT animals (two-way ANOVA,
333 $F(3,37)=0.19$, genotype*treatment interaction, $p=.906$, **Figure 5a**), indicating that 15
334 weeks of different pharmacological regimes did not differentially affect feeding
335 behavior and, possibly, feeding-related confounders on behavior. However, VMAT2-
336 LO animals appeared to eat significantly more than VMAT2-WT controls in respect to

337 their body weight (two-way ANOVA, $F(1,37)=18.36$, genotype main effect, $***p<.001$,
338 **Figure 5a**).

339

340 *3.5 Effect of chronic combined administration of L-DOPA and tolcapone on plasma* 341 *levels of tHcy and hepatic enzymes*

342 In agreement with previous evidence (Miller et al., 2003), we found that administration
343 of L-DOPA increased the levels of tHcy in plasma and pushed them over the maximum
344 standard concentration (one-way ANOVA $F(3,18)=3.49$, $*p=.037$, **Figure 5b**) when
345 compared to vehicle animals (Tukey post hoc test, $*p=.038$). As expected, vehicle and
346 tolcapone administration alone did not result in increased tHcy generation. Notably,
347 stabilizing the bioavailability of L-DOPA with tolcapone appeared to restore tHcy
348 plasma levels to normal values (Tukey post hoc test, $p=.865$, **Figure 5b**). Normal liver
349 function was assessed on the last week of treatment by measuring AST and ALT
350 plasma levels from blood samples of VMAT2-LO-V and VMAT2-LO-LT, the last being
351 considered the group with the heaviest drug regime. We found no differences in ALT
352 (unpaired $t(7)=1.49$, $p=.179$, **Figure 5c**) and AST levels (unpaired $t(7)=0.40$, $p=.703$,
353 **Figure 5c**), nor in the AST/ALT ratio (unpaired $t(7)=0.98$, $p=.359$, **Figure 5d**),
354 indicating no detrimental effect of chronic drug interventions at the assayed dosages
355 on their hepatic functionality.

356

357 **4. Discussion**

358 Whether a more continuous dopaminergic stimulation, in general, and via COMT
359 inhibition, in particular, may reduce L-DOPA-induced motor complications has been a
360 matter of debate, despite some important high-impact publications (Stocchi and

361 Olanow, 2004). Available studies in human PD patients and rodent PD models could
362 not unambiguously confirm such an effect (Katsaiti and Nixon, 2018). In particular, the
363 STRIDE-PD study applying entacapone and L-DOPA in PD patients did not reveal a
364 disease-modifying effect, but this negative result was hypothesized to be linked to
365 higher L-DOPA equivalents in the entacapone group (Stocchi et al., 2010). Thus, it is
366 conceivable that a beneficial effect from continuous dopaminergic stimulation by
367 COMT inhibition has been masked by higher L-DOPA concentrations at the receptor
368 site (Picconi et al., 2008). Apart from DA imbalance at the effector site, no studies are
369 available on the long-term histopathological effects of such a strategy, i.e. whether
370 more continuous dopaminergic stimulation via COMT inhibition reduces
371 neuropathological findings in PD subjects.

372 From a behavioral point of view, we consider the present study as negative, as the
373 main hypothesis – is adding a COMT inhibitor to pulsatile L-DOPA treatment
374 neuroprotective? – could not be fully confirmed. In this line, the improved outcome on
375 the bar test could merely reflect a prolonged efficacy of dopaminergic replacement
376 with COMT inhibition rather than a disease-modifying effect, while other behavioral
377 outcomes were similar between treatment groups. On the other hand, on an
378 immunohistochemistry level, we observed rescued DAT levels in VMAT2-LO-LT mice,
379 and no differences between vehicle- and L-DOPA- or tolcapone-treated animals. This
380 result corroborates previous work in this animal model, finding no change in DAT
381 expression after L-DOPA subchronic (what is this?) administration, although at an
382 earlier age (Reveron et al., 2002). In contrast, DAT staining [presented a statistical](#)
383 [trend towards higher DAT reactivity in VMAT2-LO-LT when compared to VMAT2-LO-](#)
384 [L was statistically similar between VMAT2-LO-L and VMAT2-LO-LT, yet this lack of a](#)
385 [difference might be explained by insufficient power.](#) Taken together, the behavioral

386 and immunohistochemical results might suggest that treating PD animals with L-
387 DOPA+tolcapone improves their functional prognosis and reduces signs of
388 neurodegeneration.

389 The main limitation of our study is the lack of assessment of multiple markers of
390 neurodegeneration upon treatments. This shortcoming is related to the treatment
391 window chosen for the study. While tyrosine hydroxylate neurons, likely the gold
392 standard marker of dopaminergic system integrity, are reported to die at a later stage
393 in VMAT2-LO animals (Taylor et al., 2011), altered synuclein deposition is found since
394 early age in this model (own unpublished data). This temporal disparity between the
395 two most conventional neurodegeneration markers in PD, rendered DAT the
396 intermediate outcome of choice. Rescued DAT levels by combined treatment,
397 however, is a pertinent outcome pointing to future reduced neurodegeneration in
398 VMAT2-LO mice. This is based not only on the fact that DAT reflects progressive
399 damage to the dopaminergic system, but also that it is fundamentally involved in the
400 profound neurochemical alterations that precede nigral degeneration in this model
401 (Miller et al., 1997; Caudle et al., 2007). The “dying back” hypothesis proposes that
402 decreased DAT function and expression might stand as a reliable proxy for loss of
403 striatal dopaminergic projections, and most importantly, as indicator of future damage
404 to the dopaminergic perikarya in the substantia nigra (Burke and O'Malley, 2013).
405 Interestingly, the deterioration of DAT expression and activity seen at the time of
406 sacrifice in VMAT2-LO-V mice is, reportedly, not resulting from loss of fiber density
407 (and concomitant cell death) nor alterations in DAT expression in the midbrain of these
408 animals (Caudle et al., 2007). ~~Lifetime intraneuronal compensatory mechanisms and~~
409 ~~interaction with DA quinones and reactive oxygen species may explain the decrease~~

~~in DAT observed in aged vehicle-treated animals (Whitehead et al., 2001).~~ Add a limitation re the VMAT2 model, see R1?

One possible read-out of this study, therefore, is that inhibiting COMT with tolcapone in conjunction with chronic L-DOPA treatment, thus preventing DA fluctuations in the brain, might reduce DAT decline and, consequently, future nigrostriatal neurodegeneration. The behavioral outcome could partially confirm this hypothesis insofar as VMAT2-LO-LT mice were the only group presenting stabilized bradykinesia and postural stability scores. Still, we cannot exclude a prolonged effect of dopaminergic stimulation at the receptor site, due to the addition of tolcapone, as discussed above. Nevertheless, the statistically significant link between behavior and immunohistochemical findings might further corroborate this hypothesis: slower deterioration of bar test performance in L-DOPA-treated groups appear correlated with preserved striatal DAT density levels, particularly the group co-administrated with the COMT inhibitor, to near the extent seen in VMAT2-WT-V group.

In relation to eventual confounders, it is worth to mention that there is no evidence of compensatory changes in VMAT2 expression (Reveron et al., 2002) and no exacerbated effect on the feeding behavior of the animals. Moreover, the present analysis was inconclusive on the tolcapone influence on tHcy levels in the transgenic animals, although tolcapone indeed reduced tHcy in VMAT2-WT-L animals. Lastly, we did not find signs of hepatotoxicity from chronic tolcapone administration, in line with early studies reporting harm of this drug only to very high administration dosages (Longo et al., 2016).

4.1 Conclusions

434 Altogether, despite equaling L-DOPA concentrations, this study does not undoubtedly
435 support the assumption that adding a central COMT inhibitor to L-DOPA therapy
436 provides a neuroprotective effect compared to L-DOPA alone. Although the results
437 here are believed to reflect the deceleration of nigrostriatal degeneration insofar the
438 integrity of the dopaminergic terminals is concerned, further experiments targeting a
439 later intervention window enclosing other functional and/or histopathologic deficits in
440 this PD model could clarify whether combining a COMT inhibitor can be
441 neuroprotective. Such an attempt would be most welcome, as the present study –
442 even as considered overall negative regarding the study hypothesis – produced results
443 which at least indicate that mildly enhanced continuous dopaminergic stimulation
444 might reduce DAT decline and, consequently, future nigrostriatal neurodegeneration.

445 From a clinical perspective, this finding would become relevant once there were strong
446 clinical evidence supporting these preclinical insights. In other words, if large clinical
447 delayed start design studies would confirm that smoothening L-DOPA fluctuations –
448 be it with COMT inhibitors or with other strategies - decelerates the disease course,
449 then clinical practice might shift towards adding a COMT inhibitor to L-DOPA plus
450 benserazide/carbidopa in early PD patients. This seems particularly relevant as
451 nowadays many neurologists shift back to initiate treatment with L-DOPA rather than
452 dopamine agonists because of the many and potentially harmful side effects (i.e.
453 impulse control disorders) of the latter.

454 **Acknowledgements:** The project was funded by the Zurich Center for Integrative
455 Human Physiology (ZIHP) and the Hurka Foundation. The authors would like to thank
456 Ms. Sedef Kollarik and Ms. Samantha Webber for their valued technical assistance.

457 **5. References**

458 Alvarez-Cervera F, Villanueva-Toledo J, Moo-Puc R, Heredia-López FJ, Alvarez-
459 Cervera M, Pineda JC, Góngora-Alfaro JL (2005) A novel automated rat catalepsy bar
460 test system based on a RISC microcontroller. *Journal of Neuroscience*
461 *Methods*, 146(1), 76-83.

462
463 Avale ME, Falzone TL, Gelman DM, Low MJ, Grandy DK, Rubinstein M (2004) The
464 dopamine D4 receptor is essential for hyperactivity and impaired behavioral inhibition
465 in a mouse model of attention deficit/hyperactivity disorder. *Molecular Psychiatry*, 9(7),
466 718-726.

467
468 Bäckström R, Honkanen E, Pippuri A, Kairisalo P, Pystynen J, Heinola K, Nissinen E,
469 Linden IB, Männistö PT, Kaakkola S, et al. (1989) Synthesis of some novel potent and
470 selective catechol O-methyltransferase inhibitors. *Journal of Medicinal*
471 *Chemistry*, 32(4), 841-846.

472
473 Bello EP, Mateo Y, Gelman DM, Noaín D, Shin JH, Low MJ, Alvarez VA, Lovinger DM,
474 Rubinstein M (2011) Cocaine supersensitivity and enhanced motivation for reward in
475 mice lacking dopamine D2 autoreceptors. *Nature Neuroscience*, 14(8), 1033-1038.

476
477 Bhattacharjee N, Borah A (2016) Oxidative stress and mitochondrial dysfunction are
478 the underlying events of dopaminergic neurodegeneration in homocysteine rat model
479 of Parkinson's disease. *Neurochem Int.*; 101:48-55.

480

Formatiert: Englisch (Vereinigtes Königreich)

481 Bibbiani F, Costantini LC, Patel R, Chase TN (2005) Continuous dopaminergic
482 stimulation reduces risk of motor complications in parkinsonian
483 primates. *Experimental Neurology*, 192(1), 73-78.

484

485 Bonifácio MJ, Palma PN, Almeida L, Soares-da-Silva P (2007) Catechol-O-
486 methyltransferase and its inhibitors in parkinson's disease. *CNS Drug Reviews*, 13(3),
487 352-379.

488

489 Bonifácio MJ, Torção L, Loureiro AI, Palma PN, Wright LC, Soares-da-Silva P (2015)
490 Pharmacological profile of opicapone, a third-generation nitrocatechol catechol-O-
491 methyl transferase inhibitor, in the rat. *British Journal of Pharmacology*, 172(7), 1739-
492 1752.

493

494 Braak H, Del Tredici K, Rüb U, de Vos RA, Jansen Steur EN, Braak E (2003). Staging
495 of brain pathology related to sporadic parkinson's disease. *Neurobiology of*
496 *Aging*, 24(2), 197-211.

497

498 Burke RE, O'Malley K (2013) Axon degeneration in Parkinson's disease. *Exp Neurol*.
499 2013 Aug;246:72-83.

500

501 Caudle WM, Richardson JR, Wang MZ, Taylor TN, Guillot TS, McCormack AL,
502 Colebrooke RE, Di Monte DA, Emson PC, Miller GW (2007) Reduced vesicular
503 storage of dopamine causes progressive nigrostriatal neurodegeneration. *The Journal*
504 *of Neuroscience*, 27(30), 8138-8148.

505

Cenci MA (2014) Presynaptic mechanisms of L-DOPA-induced dyskinesia: the findings, the debate, and the therapeutic implications. *Front. Neurol.* 5:242. doi: 10.3389/fneur.2014.00242

Chen, J., Song, J., Yuan, P., Tian, Q., Ji, Y., Ren-Patterson, R., Liu, G., Sei, Y., & Weinberger, D. R. (2011). Orientation and cellular distribution of membrane-bound catechol-O-methyltransferase in cortical neurons: implications for drug development. *The Journal of biological chemistry*, 286(40), 34752–34760. <https://doi.org/10.1074/jbc.M111.262790>

Formatiert: Englisch (Vereinigtes Königreich)

Chondrogiorgi M, Tatsioni A, Reichmann H, Konitsiotis S (2014) Dopamine agonist monotherapy in parkinson's disease and potential risk factors for dyskinesia: A meta-analysis of levodopa-controlled trials. *European Journal of Neurology*, 21(3), 433-440.

Connolly BS, Lang AE (2014) Pharmacological treatment of parkinson disease: A review. *Jama*, 311(16), 1670-1683.

Formatiert: Englisch (Vereinigtes Königreich)

~~Di Monte DA, McCormack A, Petzinger G, Janson AM, Quik M, Langston WJ (2000) Relationship among nigrostriatal denervation, parkinsonism, and dyskinesias in the MPTP primate model. *Movement Disorders*, 15(3), 459-466.~~

Giasson BI, Duda JE, Quinn SM, Zhang B, Trojanowski JQ, Lee VM (2002) Neuronal α -synucleinopathy with severe movement disorder in mice expressing A53T human α -synuclein. *Neuron*, 34(4), 521-533.

531 Guldberg HC, Marsden CA (1975) Catechol-O-methyl transferase: Pharmacological
 532 aspects and physiological role. *Pharmacol Rev*, 27(2), 135.

533

534 [Haasio, K., Koponen, A., Penttilä, K. E., & Nissinen, E. \(2002\). Effects of entacapone](#)
 535 [and tolcapone on mitochondrial membrane potential. *European journal of*](#)
 536 [pharmacology](#), 453(1), 21–26. [https://doi.org/10.1016/s0014-2999\(02\)02383-x](https://doi.org/10.1016/s0014-2999(02)02383-x)

537

538 [Ibrahimagic, O. C., Smajlovic, D., Dostovic, Z., Pasic, Z., Kunic, S., Iljazovic, A., &](#)
 539 [Hajdarevic, D. S. \(2016\). Hyperhomocysteinemia and its treatment in patients with](#)
 540 [parkinson's disease. *Materia socio-medica*, 28\(4\), 303–306.](#)
 541 <https://doi.org/10.5455/msm.2016.28.303-306>

542

543 Jenner P, McCreary AC, Scheller DKA (2011) Continuous drug delivery in early- and
 544 late-stage parkinson's disease as a strategy for avoiding dyskinesia induction and
 545 expression. *Journal of Neural Transmission*, 118(12), 1691-1702.

546

547 Kaakkola S (2000) Clinical pharmacology, therapeutic use and potential of COMT
 548 inhibitors in parkinson's disease. *Drugs*, 59(6), 1233-1250.

549

550 Katsaiti I, Nixon J (2018) Are there benefits in adding catechol-O methyltransferase
 551 inhibitors in the pharmacotherapy of parkinson's disease patients? A systematic
 552 review. *Journal of Parkinson's Disease*, 8(2), 217-231.

553

554 Kiss LE, Ferreira HS, Torráo L, Bonifácio MJ, Palma PN, Soares-da-Silva P,
 555 Learmonth DA (2010) Discovery of a long-acting, peripherally selective inhibitor of
 556 catechol-O-methyltransferase. *Journal of Medicinal Chemistry*, 53(8), 3396-3411.

557

558 [Kocer, B., Guven, H., & Comoglu, S. S. \(2016\). Homocysteine Levels in Parkinson's](#)
 559 [Disease: Is Entacapone Effective?. BioMed research international, 2016, 7563705.](#)
 560 <https://doi.org/10.1155/2016/7563705>

561

562 Kopin IJ (1985) Catecholamine metabolism: Basic aspects and clinical
 563 significance. *Pharmacol Rev*, 37(4), 333.

564

565 Kurth MC, Adler CH (1998) COMT inhibition: a new treatment strategy for Parkinson's
 566 disease. *Neurology*, 50(5 Suppl 5):S3-14.

567

568 ~~[Langston JW \(2017\) The MPTP story. *Journal of Parkinson's Disease*, 7, S11-S19.](#)~~

569

570 ~~[Leta V, Jenner P, Chaudhuri KR, Antonini A \(2019\) Can therapeutic strategies prevent](#)~~
 571 ~~[and manage dyskinesia in parkinson's disease? an update. *Expert Opinion on Drug*](#)~~
 572 ~~[Safety](#), 18(12), 1203-1218.~~

573 ~~[Lamberti, P., Zoccolella, S., Iliceto, G., Armenise, E., Fraddosio, A., De Mari, M. and](#)~~
 574 ~~[Livrea, P. \(2005\). Effects of levodopa and COMT inhibitors on plasma homocysteine](#)~~
 575 ~~[in Parkinson's disease patients. *Mov. Disord.*, 20: 69-72. doi:10.1002/mds.20261](#)~~

576

577 Longo DM, Yang Y, Watkins PB, Howel, BA, Siler SQ (2016) Elucidating differences
 578 in the hepatotoxic potential of tolcapone and entacapone with DILIsym(®), a

579 mechanistic model of drug-induced liver injury. *CPT: Pharmacometrics & Systems*
580 *Pharmacology*, 5(1), 31-39.

581

582 Miller GW, Staley JK, Heilman CJ, Perez JT, Mash DC, Rye DB, Levey AI (1997)
583 Immunochemical analysis of dopamine transporter protein in Parkinson's disease. *Ann*
584 *Neurol.*; 41(4):530-9.

585

586 Miller JW, Selhub J, Nadeau MR, Thomas CA, Feldman RG, Wolf PA (2003) Effect of
587 L-dopa on plasma homocysteine in PD patients: relationship to B-vitamin
588 status. *Neurology*. 60(7):1125–1129.

589

590 Mooslehner KA, Chan PM, Xu W, Liu L, Smadja C, Humby T, Allen ND, Wilkinson LS,
591 Emson PC (2001) Mice with very low expression of the vesicular monoamine
592 transporter 2 gene survive into adulthood: Potential mouse model for
593 parkinsonism. *Molecular and Cellular Biology*, 21(16), 5321.

594

595 Müller T (2011) Motor complications, levodopa metabolism and progression of
596 Parkinson's disease. *Expert Opin Drug Metab Toxicol.*; 7(7):847-55.

597

598 Nagatsu T, Sawada M (2009a) L-dopa therapy for parkinson's disease: Past, present,
599 and future. *Parkinsonism & Related Disorders; Proceedings of the 3rd International*
600 *Symposium on Dopaminergic and Nondopaminergic Mechanisms in Parkinson's*
601 *Disease (INPD)*, 15, S3-S8.

602

Formatiert: Englisch (Vereinigtes Königreich)

603 ~~Niceolini F, Rocchi L, Politis M (2015) Molecular imaging of levodopa-induced~~
604 ~~dyskinesias. *Cellular and Molecular Life Sciences*, 72(11), 2107-2117.~~

605

606 Noaín D, Pérez-Millán MI, Bello EP, Luque GM, Casas Cordero R, Gelman DM, Peper
607 M, Tornadu IG, Low MJ, Becú-Villalobos D, Rubinstein M (2013) Central dopamine
608 D2 receptors regulate growth-hormone-dependent body growth and pheromone
609 signaling to conspecific males. *The Journal of Neuroscience*, 33(13), 5834-5842.

610

611 Nyholm D (2006) Enteral levodopa/carbidopa gel infusion for the treatment of motor
612 fluctuations and dyskinesias in advanced parkinson's disease. *Expert Review of*
613 *Neurotherapeutics*, 6(10), 1403-1411.

614

615 Olanow CW, Schapira AHV (2013) Therapeutic prospects for parkinson
616 disease. *Annals of Neurology*, 74(3), 337-347.

617

618 Paul R, Borah A (2016) L-DOPA-induced hyperhomocysteinemia in parkinson's
619 disease: Elephant in the room. *Biochimica Et Biophysica Acta (BBA) - General*
620 *Subjects*, 1860(9), 1989-1997.

621

622 Picconi B, Paillé V, Ghiglieri V, Bagetta V, Barone I, Lindgren HS, Bernardi G, Angela
623 Cenci M, Calabresi P (2008) L-DOPA dosage is critically involved in dyskinesia via
624 loss of synaptic depotentiation. *Neurobiology of Disease*, 29(2), 327-335.

625

Formatiert: Englisch (Vereinigtes Königreich)

626 Reveron ME, Savelieva KV, Tillerson JL, McCormack AL, Di Monte DA, Miller GW
 627 (2002) L-DOPA does not cause neurotoxicity in VMAT2 heterozygote knockout mice.
 628 Neurotoxicology; 23(4-5):611-9.
 629
 630 Ruifrok AC, Johnston DA (2001) Quantification of histochemical staining by color
 631 deconvolution. *Analytical and Quantitative Cytology and Histology*, 23(4), 291-9.
 632
 633 de Ryck M, Schallert T, Teitelbaum P (1980) Morphine versus haloperidol catalepsy
 634 in the rat: A behavioral analysis of postural support mechanisms. *Brain*
 635 *Research*, 201(1), 143-172.
 636
 637 Schrag A, Quinn N (2000) Dyskinesias and motor fluctuations in Parkinson's disease.
 638 A community-based study. *Brain*. 2000, 123(Pt11), 2297–2305.
 639 Stocchi F, Olanow CW (2004) Continuous dopaminergic stimulation in early and
 640 advanced parkinson's disease. *Neurology*, 62(1), S56.
 641
 642 Schreiner SJ, Imbach LL, Werth E, Poryazova R, Baumann-Vogel H, Valko PO, Murer
 643 T, Noain D, Baumann CR (2019) Slow-wave sleep and motor progression in parkinson
 644 disease. *Annals of Neurology*, 85(5), 765-770.
 645
 646 Stocchi F, Rascol O, Kieburz K, Poewe W, Jankovic J, Tolosa E, Barone P, Lang AE,
 647 Olanow CW (2010) Initiating levodopa/carbidopa therapy with and without entacapone
 648 in early parkinson disease: The STRIDE-PD study. *Annals of Neurology*, 68(1), 18-27.
 649

650 Taylor TN, Caudle WM, Shepherd KR, Noorian A, Jackson CR, Iuvone PM,
651 Weinshenker D, Greene JG, Miller GW (2009) Nonmotor symptoms of parkinson's
652 disease revealed in an animal model with reduced monoamine storage capacity. *The*
653 *Journal of Neuroscience*, 29(25), 8103-8113.

654
655 Taylor TN, Caudle WM, Miller GW (2011) VMAT2-deficient mice display nigral and
656 extranigral pathology and motor and nonmotor symptoms of parkinson's
657 disease. *Parkinson's Disease*, 2011, 124165.

658
659 Verschuur CVM, Suwijn SR, Boel JA, Post B, Bloem BR, van Hilten JJ, van Laar T,
660 Tissingh G, Muntz AG, Deuschl G, Lang AE, Dijkgraaf MGW, de Haan RJ, de Bie
661 RMA; LEAP Study Group (2019) Randomized delayed-start trial of levodopa in
662 parkinson's disease. *N Engl J Med*, 380(4), 315-324.

663
664 Whitehead RE, Ferrer JV, Javitch JA, Justice JB (2001) Reaction of oxidized
665 dopamine with endogenous cysteine residues in the human dopamine transporter. *J*
666 *Neurochem*. 76(4):1242–1251.

667
668 ~~Winkler C, Kirik D, Björklund A, Cenci MA (2002) L-DOPA-induced dyskinesia in the~~
669 ~~intrastratial 6-hydroxydopamine model of parkinson's disease: Relation to motor and~~
670 ~~cellular parameters of nigrostriatal function. *Neurobiology of Disease*, 10(2), 165-186.~~

671
672 Zeuner KE, Schäffer E, Hopfner F, Brüggemann N, Berg D (2019) Progress of
673 Pharmacological Approaches in Parkinson's Disease. *Clin Pharmacol Ther.*,
674 105(5):1106-1120.

Formatiert: Italienisch (Schweiz)

675
676 [Zoccolella S, Martino D, Defazio G, Lamberti P, Livrea P. \(2006a\).](#)
677 [Hyperhomocysteinemia in movement disorders: Current evidence and hypotheses.](#)
678 [Curr Vasc Pharmacol.;4\(3\):237-243. doi:10.2174/157016106777698414](#)
679
680 [Zoccolella, S., Lamberti, P., Iliceto, G., Dell'Aquila, C., Diroma, C., Fraddosio, A.,](#)
681 [Lamberti, S. V., Armenise, E., Defazio, G., Mari, M., & Livrea, P. \(2006b\). Elevated](#)
682 [plasma homocysteine levels in L-dopa-treated Parkinson's disease patients with](#)
683 [dyskinesias, Clinical Chemistry and Laboratory Medicine \(CCLM\), 44\(7\), 863-866. doi:](#)
684 <https://doi.org/10.1515/CCLM.2006.143>
685
686 Zürcher G, Colzi A, Da Prada M (1990) Ro 40-7592: Inhibition of COMT in rat brain
687 and extracerebral tissues. Paper presented at the *Amine Oxidases and their Impact*
688 *On*, 375-380.

689 **Figure legends**

690

691 **Figure 1. Experimental design and timeline.** VMAT2 LO (n = 21) and WT (n = 23)
692 mice were behaviorally tested at 14 months of age (baseline: BL), before starting a
693 chronic regime of pharmacological interventions that lasted 17 weeks. Each treatment
694 month consisted of 3 weeks of drug administration ('ON'-medication, blue rectangles),
695 plus 1 week of behavioral testing ('OFF'-medication, grey rectangle). Each ON-
696 medication week consisted of 5 days of twice daily per os dosaging, at light onset and
697 8 hours late, of saline (-V), L-DOPA (-L), tocapone (-T) or L-DOPA+tolcapone (-LT).
698 On the 15th ON-medication week, the food intake was assessed in all animals. After
699 17 weeks on interventions, the mice were sacrificed (red marking) and blood and
700 tissue samples collected for posterior evaluations. (AST – aspartate transaminase,
701 ALT – alanine transaminase, tHcy – total homocysteine, DAT – dopamine transporter).

702

703 **Figure 2. Motor ability in response to chronic pharmacological interventions in**
704 **VMAT2-LO mice.** At baseline (BL), VMAT2-LO (n = 21) and VMAT2-WT control (n =
705 23) drug naïve mice were behaviorally tested in 3 different motor domains: **(a)**
706 locomotion (open field test), **(b)** body rigidity (bar test), and **(c)** coordination (rotarod
707 test). In all three tests, VMAT2-LO mice performed significantly worse than VMAT2-
708 WT **((a)** unpaired $t(42)=6.03$, $***p<.001$; **(b)** Welch's $t(20)=4.56$, $***p<.001$; **(c)**
709 unpaired $t(42)=4.17$, $***p<.001$). At endpoint testing, after 16 weeks of interventions,
710 all VMAT2-LO mice presented a marked behavioral impairment in all parameters
711 explored, as compared to VMAT2-WT controls **((d)** nested 2-way ANOVA,
712 treatment(genotype), $F(4,36)=10.43$, $***p<.001$; **(e)** nested 2-way ANOVA of ranked
713 cases, treatment(genotype), $F(4,36)=23.06$, $***p<.001$; **(f)** nested 2-way ANOVA of

714 ranked cases, treatment(genotype), $F(4,36)=34.45$, $***p<.001$). The progression of
 715 individual symptoms in VMAT2-LO mice in respect to the treatment received was
 716 analyzed in subsets: (g, h, i) -V vs -T treated VMAT2-LO animals were compared to
 717 evidence a potential stand-alone effect of the drug, aside from its effect over L-DOPA
 718 stability; (j, k, l) -L vs -LT treated VMAT2-LO animals were compared to evidence the
 719 effect of stabilizing L-DOPA bioavailability on motor progression. -V vs -T progressions
 720 were not significantly different (t-test on grouped slopes), while -L vs -LT progressions
 721 in the bar test differed significantly (k, unpaired $t(8.43)=4.24$, $**p=.003$), with the latter
 722 evidencing practically no worsening of body rigidity in VMAT2-LO (slope of dark line
 723 close to 1) over a 17-week period. (V = vehicle; L = L-DOPA; T = tolcapone; LT = L-
 724 DOPA+tolcapone; number of animals: WT-V=5, WT-L=6, WT-T=6, WT-LT=6, LO-
 725 V=5, LO-L=5, LO-T=5, LO-LT=6).

Formatiert: Italienisch (Schweiz)

727 **Figure 3. Preserved striatal DAT integrity in VMAT2-LO with stabilized L-DOPA**
 728 **bioavailability.** (a) Analyzed area from coronal sections of striatum from VMAT2-WT
 729 and VMAT2-LO mice. (b) VMAT2-LO-V mice presented reduced DAT immunostaining
 730 strength compared to VMAT2-WT-V controls (Welch's $t(29.86)=2.43$, $*p=.022$). (c)
 731 Individual examples of DAT reactivity in striatum of -V, -L, -T and -LT VMAT2-LO
 732 animals. (d) Statistical analyses of optical-density in differently treated VMAT2-LO
 733 animals show partially rescued DAT levels in VMAT2-LO-LT (one-way ANOVA,
 734 Brown-Forsythe $F(3,69)=4.83$, $V*LT$, $**p=.004$). (V = vehicle; L = L-DOPA; T =
 735 tolcapone; LT = L-DOPA+tolcapone; number of animals: WT-V=5, LO-V=3, LO-L=5,
 736 LO-T=5, LO-LT=6).

737 **Figure 4. Rigidity correlates with integrity of striatal dopaminergic terminals. (a)**
 738 The slope of the robust linear fit on the time of immobility' scores of each VMAT2-LO
 739 individual over the 17-week treatment, representing the progression of bradykinesia
 740 during the complete intervention (<0 = improvement, >0 = worsening), and DAT optical
 741 density, as a marker of DA striatal terminal integrity, in VMAT2-LO mice correlate
 742 significantly (Pearson's, $r(17)=-0.50$, $*p=.015$, $n = 19$). (b) Exploratory group-based
 743 clustering for bradykinesia progression (<0 means improvement, >0 means
 744 worsening) and DAT optical density levels for VMAT2-WT-V and VMAT2-LO-V, -L, -T
 745 and -LT groups. (V = vehicle; L = L-DOPA; T = tolcapone; LT = L-DOPA+tolcapone;
 746 number of animals: WT-V=5, LO-V=3, LO-L=5, LO-T=5, LO-LT=6).

747
 748 **Figure 5. Effects on food intake, tHcy plasma levels and hepatic enzymes. (a)**
 749 Food intake was evaluated on protocol-week 15 for VMAT2-LO ($n = 21$) and VMAT2-
 750 WT ($n = 23$) mice. Ingested food per body weight was calculated for each group: no
 751 differences between treatments within genotypes were observed (two-way ANOVA,
 752 $F(3,37)=0.19$, genotype*treatment interaction, $p=.906$). However, when pulling
 753 together all animals of the same genotype, regardless of the treatment, VMAT2-LO
 754 animals show increased feeding behavior (two-way ANOVA, $F(1,37)=18.36$, genotype
 755 main effect, $***p<.001$) when compared to VMAT2-WT mice. Number of animals: WT-
 756 V=5, WT-L=6, WT-T=6, WT-LT=6, LO-V=5, LO-L=5, LO-T=5, LO-LT=6. (b) Increased
 757 tHcy plasma levels observed upon L-DOPA treatment were normalized by combined
 758 L-DOPA+tolcapone (one-way ANOVA $F(3,18)=3.49$, $*p=.037$; Tuckey multiple-
 759 comparisons: V*L, $*p=.038$; V*LT, $p=.865$). Shadow indicates normal plasma tHcy
 760 levels ($< 6 \mu\text{mol/L}$) in male C57BL/6J mice. Number of animals: WT-V=5, WT-L=6,
 761 WT-T=6, WT-LT=6. (c) Alanine transaminase (ALT) and aspartate transaminase

762 (AST), marker enzymes of functional liver integrity, were determined in -V and -LT
763 VMAT2-LO mice. No differences were observed between treatments (ALT, V*LT,
764 unpaired $t(7)=1.49$, $p=.179$; AST, V*LT, unpaired $t(7)=0.40$, $p=.703$). Number of
765 animals: LO-V=5, LO-LT=4. (d) The ratio of AST/ALT was determined to also be
766 normal (AST/ALT ratio, V*LT, unpaired $t(7)=0.98$, $p=.359$). V = vehicle; L = L-DOPA;
767 T = tolcapone; LT = L-DOPA+tolcapone.